

International Journal of Modern Physics E
 © World Scientific Publishing Company

TRITON-³HE RELATIVE AND DIFFERENTIAL FLOWS AND THE HIGH DENSITY BEHAVIOR OF NUCLEAR SYMMETRY ENERGY

GAO-CHAN YONG

*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, P.R. China
 yonggaochan@impcas.ac.cn*

BAO-AN LI

*Department of Physics and Astronomy, Texas A&M University-Commerce, Commerce, TX
 75429-3011, USA
 Bao-An.Li@tamuc-commerce.edu*

LIE-WEN CHEN

*Department of Physics, Shanghai Jiao Tong University, Shanghai 200240, China
 Center of Theoretical Nuclear Physics, National Laboratory of Heavy-Ion Accelerator, Lanzhou,
 730000, China
 lwchen@sjtu.edu.cn*

Received (received date)

Revised (revised date)

Using a transport model coupled with a phase-space coalescence after-burner we study the triton-³He relative and differential transverse flows in semi-central ¹³²Sn + ¹²⁴Sn reactions at a beam energy of 400 MeV/nucleon. We find that the triton-³He pairs carry interesting information about the density dependence of the nuclear symmetry energy. The t-³He relative flow can be used as a particularly powerful probe of the high-density behavior of the nuclear symmetry energy.

1. Introduction

The density dependence of nuclear symmetry energy especially at supra-saturation densities is among the most uncertain properties of neutron-rich nuclear matter^{1,2}. However, it is very important for nuclear structure^{3,4}, heavy-ion reactions^{5,6,7,8,9,10} and many phenomena/processes in astrophysics and cosmology^{11,12,13}. Heavy-ion reactions especially those induced by radioactive beams provide a unique opportunity to constrain the symmetry energy at supra-saturation densities in terrestrial laboratories. Various probes using heavy-ion reactions have been proposed in the literature, see, e.g., ref.¹⁰ for the most recent review. It is particularly interesting to mention that, besides many significant results about the symmetry energy at sub-saturation densities, see, e.g., refs.^{14,15,16,17,18,19,20}, circumstantial evidence for a rather soft symmetry energy at supra-saturation densities has been reported very recently²¹ based on the IBUU04 transport model²² analysis

of the π^-/π^+ data taken by the FOPI Collaboration at SIS/GSI²³. To constrain tightly and reliably the nuclear symmetry energy especially at supra-saturation densities, much more efforts by both the nuclear physics and the astrophysics communities are still needed.

There is an urgent need to verify the conclusion about the soft symmetry energy at supra-saturation densities required to reproduce the FOPI π^-/π^+ data within transport model analyses^{21,23}. It was predicted that the neutron-proton differential flow is another sensitive probe of the high-density behavior of the nuclear symmetry energy²⁴. However, it is difficult to measure observables involving neutrons. One question often asked by some experimentalists is whether the triton-³He pair may carry the same information as the neutron-proton one. We will try to answer this question quantitatively by coupling the IBUU04 calculations to a phase-space coalescence after-burner. Indeed, we found that, similar to the neutron-proton pair, the triton-³He relative and differential transverse flows are sensitive to the high-density behavior of the nuclear symmetry energy. They can be used to test indications about the high-density behavior of the symmetry energy observed earlier from analyzing the π^-/π^+ data.

2. The theoretical models

Our study is carried out based on the IBUU04 version of an isospin and momentum dependent transport model and the simplest phase-space coalescence after-burner. The single nucleon potential is one of the most important inputs to BUU-like transport models for nuclear reactions. In the IBUU04 transport model, we use a single nucleon potential derived within the Hartree-Fock approach using a modified Gogny effective interaction (MDI)^{22,25}. The corresponding MDI symmetry energy can be found in Ref.²⁶

Because most BUU-type transport models including the IBUU04 are incapable of forming dynamically realistic nuclear fragments, some types of after-burners, such as statistical and coalescence models, are normally used as a remedy. This kind of hybrid models can be used to study reasonably well, for instance, nuclear multifragmentation, see, e.g., ref.^{27,28,29,30}, collective flow of light fragments, see e.g.,^{31,32,33} and the formation of hypernuclei³⁴. There are, however, some remaining issues, such as the freeze-out time of fragments that is related to the time of coupling the transport model with the after-burner, etc. There are also interesting work in using advanced coalescence models^{35,36}, see, e.g., refs.^{37,38}. We notice here that, several advanced cluster recognition routines, such as, the Early Cluster Recognition Algorithm (ECRA)³⁹, the Simulated Annealing Clusterization Algorithm (SACA)⁴⁰, have been put forward in recent years. For the purposes of the present exploration, however, we use the simplest phase-space coalescence model, see, e.g., refs.^{32,33}, where a physical fragment is formed as a cluster of nucleons with relative momenta smaller than P_0 and relative distances smaller than R_0 . The results presented in the following are obtained with $P_0 = 263$ MeV/c and $R_0 = 3$ fm.

This simple choice may thus limit the scope and importance of our study here. For instance, we shall limit ourselves to studies of the relative/differential observables for t - ^3He pairs without attempting to study pairs of the heavier mirror nuclei. An extended study including the heavy mirror nuclei using the advanced coalescence and/or earlier cluster recognition methods is planned.

3. Results and discussions




Fig. 1. Triton and ^3He transverse flows (in unit of MeV) as functions of the reduced C.M. rapidity in the semi-central $^{132}\text{Sn} + ^{124}\text{Sn}$ reactions at a beam energy of 400 MeV/nucleon. Taken from Ref. 41.

We now investigate whether the transverse collective flows of triton and ^3He can be used to probe the symmetry energy. Firstly, we examine in Fig. 1 their transverse flows individually. The average C.M. transverse momentum per nucleon $\langle p_x/A \rangle$ in the reaction plane is defined as

$$\langle p_x/A \rangle(y) \equiv \frac{1}{N(y)} \sum_{i=1}^{N(y)} p_x^i/A(y) \quad (1)$$

where $N(y)$ is the total number of fragments of mass A in the rapidity bin at y . The correlation between the $\langle p_x/A \rangle$ and rapidity y reveals the transverse collective flow⁴². It is seen that ^3He clusters show a stronger flow than triton clusters. This is mainly due to the stronger Coulomb force experienced by the ^3He clusters. More interestingly, the transverse flow of ^3He clusters show appreciable sensitivity to the variation of the symmetry energy.

The transverse flow is a result of actions of several factors including the isoscalar, symmetry and Coulomb potentials and nucleon-nucleon scatterings. It is well known




Fig. 2. Triton- ^3He relative and differential flows (in unit of MeV) as a function of the reduced C.M. rapidity in the same reaction as in Fig. 1. Taken from Ref. 41.

that the transverse flow is sensitive to the isoscalar potential. Given the remaining uncertainties associated with the isoscalar potential and the small size of the symmetry energy effects, it would be very difficult to extract any reliable information about the symmetry energy from the individual flows of triton and ^3He clusters. Thus techniques of reducing effects of the isoscalar potential while enhancing effects of the isovector potential are very helpful^{24,43,44,45,46}. We thus study in Fig. 2 the triton- ^3He relative and differential flows. The relative flow is given as

$$\langle p_x^t/A \rangle - \langle p_x^{3\text{He}}/A \rangle = \frac{1}{N_t} \sum_{i=1}^{N_t} p_x^i/A - \frac{1}{N_{3\text{He}}} \sum_{i=1}^{N_{3\text{He}}} p_x^i/A. \quad (2)$$

The triton- ^3He differential flow reads

$$\begin{aligned} \langle p_x^{t-3\text{He}}/A \rangle &= \frac{1}{N_t + N_{3\text{He}}} \left(\sum_{i=1}^{N_t} p_x^i/A - \sum_{i=1}^{N_{3\text{He}}} p_x^i/A \right) \\ &= \frac{N_t}{N_t + N_{3\text{He}}} \langle p_x^t/A \rangle - \frac{N_{3\text{He}}}{N_t + N_{3\text{He}}} \langle p_x^{3\text{He}}/A \rangle, \end{aligned} \quad (3)$$

where N_t , $N_{3\text{He}}$ are the number of triton and ^3He in the rapidity bin at y . From the upper panel of Fig. 2, it is seen that the triton- ^3He relative flow is very sensitive to the symmetry energy. Because of the larger slope of the ^3He flow, the triton- ^3He relative flow shows a negative slope at mid-rapidity. Effects of the symmetry

energy on the differential flow shown in the lower panel, however, is relatively small. Although the ³He flow is more sensitive to the symmetry energy, the small number of ³He clusters makes the ³He flow contributes less to the triton-³He differential flow (as indicated in Eq. (3)). The triton-³He differential flow is therefore dominated by triton clusters. Consequently, it is less sensitive to the symmetry energy than the triton-³He relative flow. The slope $F(x) \equiv d \langle p_x/A \rangle / d(y/y_{beam})$ of the transverse flow at mid-rapidity can be used to characterize more quantitatively the symmetry energy effects. We found that for the t-³He relative flow, $F(x=1) \approx -74$ MeV/c and $F(x=-1) \approx -22$ MeV/c, respectively. For the t-³He differential flow, $F(x=1) \approx 21$ MeV/c and $F(x=-1) \approx 42$ MeV/c, respectively. The t-³He relative flow thus can be used as a very useful and independent tool to test the soft symmetry energy at supra-saturation densities extracted from studying the π^-/π^+ ratio²¹.

4. Summary

In summary, using a hybrid approach coupling the transport model IBUU04 to a phase-space coalescence after-burner we studied the t-³He relative and differential flows in semi-central ¹³²Sn + ¹²⁴Sn reactions at an incident energy of 400 MeV/nucleon. We found that the nuclear symmetry energy affects strongly the t-³He relative and differential flows. The t-³He relative flow can be used as a particular powerful probe of the high-density behavior of the nuclear symmetry energy.

5. Acknowledgements

This work was supported in part by the US National Science Foundation Awards PHY-0652548 and PHY-0757839, the Research Corporation under Award No.7123 and the Texas Coordinating Board of Higher Education Award No.003565-0004-2007, the National Natural Science Foundation of China under grants 10710172, 10575119, 10675082 and 10975097 and MOE of China under project NCET-05-0392, Shanghai Rising-Star Program under Grant No.06QA14024, the SRF for ROCS, SEM of China, and the National Basic Research Program of China (973 Program) under Contract No.2007CB815004.

References

1. M. Kutschera, Phys. Lett. **B340**, 1 (1994).
2. S. Kubis and M. Kutschera, Acta Phys. Pol. **B30**, 2747 (1999); Nucl. Phys. **A720**, 189 (2003).
3. B.A. Brown, Phys. Rev. Lett. **85**, 5296 (2000).
4. C. J. Horowitz and J. Piekarewicz, Phys. Rev. Lett. **86**, 5647 (2001).
5. B.A. Li, C.M. Ko and W. Bauer, Int. Jour. Mod. Phys. E **7**, 147 (1998).
6. Isospin Physics in Heavy-Ion Collisions at Intermediate Energies, Eds. Bao-An Li and W. Udo Schröder (Nova Science Publishers, Inc, New York, 2001).
7. P. Danielewicz, R. Lacey, W.G. Lynch, Science **298**, 1592 (2002).

8. V. Baran, M. Colonna, V. Greco and M. Di Toro, *Phys. Rep.* **410**, 335 (2005).
9. L.W. Chen, C.M. Ko, B.A. Li, and G.C. Yong, *Front. Phys. China* **2**, 327 (2007).
10. B.A. Li, L.W. Chen and C.M. Ko, *Phys. Rep.* **464**, 113 (2008).
11. K. Sumiyoshi and H. Toki, *Astrophys. J.* **422**, 700 (1994).
12. J.M. Lattimer and M. Prakash, *Science* **304**, 536 (2004).
13. A.W. Steiner et al., *Phys. Rep.* **411**, 325 (2005).
14. L.W. Chen, C.M. Ko, and B.A. Li, *Phys. Rev. Lett.* **94**, 032701 (2005).
15. B.A. Li and L.W. Chen, *Phys. Rev.* **C72**, 064611 (2005).
16. L.W. Chen, C.M. Ko, and B.A. Li, *Phys. Rev.* **C72**, 064309 (2005) [arXiv:nucl-th/0509009].
17. D. Shetty, S.J. Yennello, and G.A. Souliotis, *Phys. Rev.* **C75**, 034602 (2007).
18. M. B. Tsang, Yingxun Zhang, P. Danielewicz, M. Famiano, Zhuxia Li, W. G. Lynch, and A. W. Steiner, *Phys. Rev. Lett.* **102**, 122701 (2009).
19. M. Centelles, X. Roca-Maza, X. Vias, and M. Warda, *Phys. Rev. Lett.* **102**, 122502 (2009).
20. G. Lehaut, F. Gulminelli, and O. Lopez, *Phys. Rev. Lett.* **102**, 142503 (2009).
21. Z.G. Xiao, B.A. Li, L.W. Chen, G.C. Yong, and M. Zhang, *Phys. Rev. Lett.* **102** 062502 (2009); M. Zhang, Z.G.Xiao, B.A. Li, L.W. Chen, G.C. Yong, and S.J. Zhun, *Phys. Rev.* **C80**, 034616 (2009).
22. B.A. Li, C.B. Das, S. Das Gupta, and C. Gale, *Nucl. Phys.* **A735**, 563 (2004); *Phys. Rev.* **C69**, 064602 (2004).
23. W. Reisdorf et al. for the FOPI Collaboration, *Nucl. Phys. A* **781**, 459 (2007).
24. B.A. Li, *Phys. Rev. Lett.* **85**, 4221 (2000).
25. C. B. Das, S. Das Gupta, C. Gale, and B.A. Li, *Phys. Rev.* **C67**, 034611 (2003).
26. J. Xu, L. W. Chen, B. A. Li and H. R. Ma, *Astrophys. J.* 697, 1549 (2009).
27. H. Kruse, B.V. Jacak, J.J. Molitoris, G.D. Westfall, and H. Stöcker, *Phys. Rev.* **C31** (1985) 1770.
28. B.A. Li, A.R. DeAngelis, and D.H.E. Gross, *Phys. Lett.* **B303**, 225 (1993).
29. K. Hagel et al., *Phys. Rev.* **C62**, 034607 (2000).
30. W.P. Tan et al., *Phys. Rev.* **C64**, R051901 (2001).
31. V. Koch et al., *Phys. Lett.* **B241**, 174 (1990).
32. L.W. Chen, F.S. Zhang, and G.M. Jin, *Phys. Rev.* **C58**, 2283 (1998).
33. F.S. Zhang, L.W. Chen, Z.Y. Ming, and Z.Y. Zhu, *Phys. Rev.* **C60**, 064604 (1999).
34. T. Gaitanos, H. Lenske, U. Mosel, *Phys. Lett.* **B663**, 197 (2008).
35. R. Mattiello et al., *Phys. Rev. Lett.* **74** 2180 (1995); R. Mattiello et al., *Phys. Rev.* **C 55**, 1443 (1997).
36. R. Scheibl, Ulrich Heinz, *Phys. Rev.* **C59**, 1585 (1999).
37. L.W. Chen, C.M. Ko, and B.A. Li, *Phys. Rev.* **C68**, 017601 (2003); *Nucl. Phys.* **A729**, 809 (2003).
38. L.W. Chen, C.M. Ko, and B.A. Li, *Phys. Rev.* **C69**, 054606 (2004).
39. A. Strachan and C. O. Dorso, *Phys. Rev.* **C56**, 995 (1997).
40. Rajeev K. Puri and Joerg Aichelin, *J. Comput. Phys.* **162**, 245 (2000).
41. G.C. Yong, B.A. Li, L.W. Chen, and X.C. Zhang, *Phys. Rev.* **C80**, 044608 (2009).
42. P. Danielewicz and G. Odyniec, *Phys. Lett.* **B157**, 146 (1985).
43. G.C. Yong, B.A. Li, and L.W. Chen, *Phys. Rev.* **C74**, 064617 (2006); *Phys. Lett.* **B650**, 344 (2007).
44. B.A. Li, L.W. Chen, G. C. Yong, and W. Zuo, *Phys. Lett.* **B634**, 378 (2006).
45. V. Greco, V. Baran, M. Colonna, M. Di Toro, T. Gaitanos, and H.H. Wolter, *Phys. Lett.* **B562**, 215 (2003).
46. M.A. Famiano, et al., *Phys. Rev. Lett.* **97**, 052701 (2006).